

**THE MOUNT SINAI HOSPITAL
2020 ADULT CUMULATIVE INPATIENT GRAM-POSITIVE ISOLATES (PERCENT SUSCEPTIBLE OF INITIAL ISOLATES)**

ADULT INPATIENT GRAM-POSITIVE		Beta-Lactams					Other Mechanisms of Action									
		Ceftriaxone, meningitis	Ceftriaxone, non-meningitis	Ampicillin	Penicillin IV, meningitis	Penicillin IV, non-meningitis	Oxacillin ^a	Clindamycin	Daptomycin	Linezolid	Doxycycline	Nitrofurantoin ^b	Trimethoprim- sulfamethoxazole	Vancomycin	Levofloxacin	Gentamicin ^c
<i>n</i>																
<i>Enterococcus faecalis</i>	359			96					99	98	23	97		89		67
<i>Staphylococcus aureus</i> MSSA	319						100	66	100	100	99 ^d		99	100		99
<i>Staphylococcus aureus</i> MRSA	257							59	100	100	96 ^d		94	100		89
<i>Enterococcus faecium</i>	253			8					85	98	38 ^d	51		21		66
<i>Staphylococcus epidermidis</i>	144						<u>16</u>	43			82 ^d			100		68
<i>Streptococcus pneumoniae</i> ^e	39	92	98		52	96								100	97	

Underlined values represent a 10% or greater decrease in susceptibility compared with 2019

a Cephalosporin (e.g., cefazolin) activity inferred by activity of oxacillin for *Staphylococcus* spp.

b Urinary tract isolates only

c Used for synergistic purposes only

d Percent susceptible based on two years of data combined (2019 and 2020)

e Percent susceptible based on three years of data combined (2018, 2019, and 2020)